

Improvement of the hot QCD pressure by the minimal sensitivity criterion

Masahiro INUI^{*)}, Akira NIÉGAWA^{**)} and Hideaki OZAKI^{***)}

Graduate School of Science, Osaka City University, Osaka 558-8585, Japan

The principles of minimal sensitivity (PMS) criterion is applied to the perturbative free energy density, or pressure, of hot QCD, which include the $\sim g_s^6 \ln g_s$ and part of the $\sim g_s^6$ terms. Applications are made separately to the short- and long-distance parts of the pressure. Comparison with the lattice results, at low temperatures, shows that the resultant “optimal” approximants are substantially improved when compared to the $\overline{\text{MS}}$ results. In particular, for the realistic case of three quark flavors, the “optimal” approximants are comparable with the lattice results.

§1. Introduction

In statistical physics, the free energy density, or pressure, plays a central role. At high temperature, the perturbative expansion of the pressure of the quark-gluon plasma is known to fifth order¹⁾ in g_s (the QCD coupling constant) and partially to sixth order.²⁾ These results are reliable only when g_s is sufficiently small. Because of the asymptotic freedom of QCD, this is achieved at high temperature $T \gtrsim 2 \text{ GeV}$ which is order of magnitude larger than the critical temperature (T_c) of the QCD phase transition. For $T_c \lesssim T \lesssim 2 \text{ GeV}$, the accuracy of the approximation, within the $\overline{\text{MS}}$ scheme, does not improve by the inclusion of higher order terms (convergence issue). Moreover, the unphysical dependence on the renormalization scale μ is strong. With the traditional choice $\mu = 2\pi T$, the truncated perturbation series for the pressure, in the $\overline{\text{MS}}$ scheme, does not reconcile with the lattice results ($0.3 \text{ GeV} \lesssim T \lesssim 0.5 \text{ GeV}$). To get rid of these dilemmas, various kinds of resummations are proposed. Among those are quasi-particle models,³⁾ Φ -derivative approximations,⁴⁾ screened perturbation theory,^{5),6)} hard thermal loop perturbation theory,⁷⁾ optimizations based on the principles of minimal sensitivity (PMS) and the fastest apparent convergence (FAC) criterions,⁶⁾ and the Padé type resummation and its variants.^{8),9)}

In this paper, putting aside the convergence issue, we will obtain^{†)} the “optimal” approximants for the pressure p through an application of the PMS criterion.¹⁰⁾ Throughout this paper, for the expression p (in massless QCD), we use the one

^{*)} E-mail: inui@sci.osaka-cu.ac.jp

^{**)} E-mail: niegawa@sci.osaka-cu.ac.jp

^{***)} E-mail: hozaki@sci.osaka-cu.ac.jp

^{†)} Blaizot, Iancu, and Rebhan were the first who, among other things, applied the PMS criterion to the perturbative pressure.⁶⁾ The content of this (PMS) part of Ref. 6) and our way of improving it will be mentioned as occasions arise later.

given^{*)} in Ref. 9):

$$p = p_E + p_M + p_G ,$$

where p_E , p_M , and p_G come, in respective order, from the hard ($\sim T$), soft ($\sim g_s T$), and supersoft ($\sim g_s^2 T$) energy regions. The scales $g_s T$ and $g_s^2 T$ are the color-electric and color-magnetic screening scales, respectively. A crucial observation made in Ref. 9) is that the “short-distance pressure” p_E and the “long-distance pressure” p_{M+G} ($\equiv p_M + p_G$) are *separately physical quantities*, and thus both, when calculated to all orders of perturbation theory, are independent of the renormalization scheme. Then we will apply the PMS criterion to p_E and p_{M+G} separately^{**)} .

In §2, a brief review of the PMS criterion is given. Then, we present the perturbation expansion of p_E and p_{M+G} in the form given in Ref. 9), which is taken as the basis of our analysis. In §3, we apply the PMS criterion to p_E and p_{M+G} separately and find the optimal approximants. In §4, we compare our results with the lattice results and the $\overline{\text{MS}}$ results. §5 is devoted to discussion and conclusion.

§2. Preliminaries

2.1. Brief review of the Principles of Minimal Sensitivity (PMS) (Ref. 10))

The couplant $\alpha_s = g_s^2/(4\pi)$ in massless QCD runs according to the renormalization group equation,

$$\mu \frac{\partial \alpha_s}{\partial \mu} = \beta(\alpha_s) = -\alpha_s^2 \sum_{j=0}^{\infty} \beta_j \alpha_s^j . \quad (2.1)$$

Here μ is the renormalization scale and

$$\begin{aligned} \beta_0 &= \frac{1}{2\pi} \left(11 - \frac{2}{3} n_f \right) , \\ \beta_1 &= \frac{1}{4\pi^2} \left(51 - \frac{19}{3} n_f \right) , \end{aligned} \quad (2.2)$$

with n_f is number of quark flavors. The renormalization schemes can be labelled by a countable infinity of parameters $(\mu, \beta_2, \beta_3, \dots)$. [β_0 and β_1 are independent of the renormalization scheme.] In the $\overline{\text{MS}}$ scheme,

$$(\beta_2)_{\overline{\text{MS}}} = \frac{1}{64\pi^3} \left(2857 - \frac{5033}{9} n_f + \frac{325}{27} n_f^2 \right) . \quad (2.3)$$

The dependence of α_s on β_j ($j = 2, 3, \dots$) is governed by

$$\frac{\partial \alpha_s}{\partial \beta_j} = -\beta(\alpha_s) \int_0^{\alpha_s} \frac{x^{j+2}}{(\beta(x))^2} dx \quad (\equiv \gamma_j(\alpha_s)) , \quad (2.4)$$

^{*)} How to construct it from the available calculations are elucidated in Ref. 9) (see also Ref. 6)).

^{**)} Since little information is available on p_G , we will apply the PMS criterion to p_{M+G} and not separately to p_M and p_G . Incidentally, in Ref. 6), the PMS criterion is applied to the total pressure $p = p_E + p_{M+G}$.

$$= \frac{2\pi}{9(j-1)}\alpha_s^{j+1} + \dots \quad (j = 2, 3, \dots). \quad (2.5)$$

It is to be noted that a change in μ results in a change in α_s of $O(\alpha_s^2)$ while a change in β_i ($i = 2, 3, \dots$) results in a change in α_s of $O(\alpha_s^{i+1})$. The running $\alpha_s = \alpha_s(\mu, \beta_2, \beta_3, \dots)$ is obtained by solving Eqs. (2.1) and (2.4) under the boundary condition that is provided by the experiment.

Let $\mathcal{R}$ be some physical quantity, $\mathcal{R} = \alpha_s^l \sum_{j=0}^{\infty} r_j \alpha_s^j$. Then, when calculated to all orders of perturbation theory, $\mathcal{R}$ is independent of the renormalization scheme, or, equivalently, of the parameters $\mu, \beta_2, \beta_3, \dots$.

The i -th order perturbative approximation $\mathcal{R}^{(i)}$ of $\mathcal{R}$ is defined by (i) truncating the perturbation series for $\mathcal{R}$ at i th order, and (ii) replacing the expansion parameter α_s by its i th-order approximant defined as the solution to Eqs. (2.1) and (2.4) with β on the right-hand sides (RHS's) truncated at i th order;

$$\mathcal{R}^{(i)} = \alpha_s^l \sum_{j=0}^{i-1} r_j \alpha_s^j, \quad \beta^{(i)}(\alpha_s) = -\alpha_s^2 \sum_{j=0}^{i-1} \beta_j \alpha_s^j, \quad (2.6)$$

$$\mu \frac{\partial \alpha_s}{\partial \mu} = \beta^{(i)}(\alpha_s), \quad \frac{\partial \alpha_s}{\partial \beta_j} = \gamma_j^{(i)}(\alpha_s), \quad (2.7)$$

where $\gamma_j^{(i)}(\alpha_s)$ is defined by Eq. (2.4) with $\beta^{(i)}$ for β .

The PMS criterion¹⁰⁾ for obtaining the optimal approximant for $\mathcal{R}$ goes as follows.

- 1) Compute $\beta^{(i)}$ and $\mathcal{R}^{(i)}$ in a particular renormalization scheme, say, $\overline{\text{MS}}$ scheme.
- 2) Impose the requirement,

$$\mu \frac{\partial \mathcal{R}^{(i)}}{\partial \mu} = \frac{\partial \mathcal{R}^{(i)}}{\partial \beta_j} = 0 \quad (j = 2, 3, \dots, i-1). \quad (2.8)$$

In the following we refer Eqs. (2.8) and (2.7) to as the OPT equations. The OPT equations determine first (a) the dependence of the coefficients ($r_1, \dots, r_{i-1}$) in Eq. (2.6) on $(\mu, \beta_2, \dots, \beta_{i-1})$, and then (b) the values for the parameters $(\mu, \beta_2, \dots, \beta_{i-1}) = (\mu^{(i)}, \beta_2^{(i)}, \dots, \beta_{i-1}^{(i)})$ together with $\alpha_s = \alpha_s^{(i)}$. From (a) and (b), the values for the coefficients are determined, $(r_1, \dots, r_{i-1}) = (r_1^{(i)}, \dots, r_{i-1}^{(i)})$.

- 3) The optimal approximant for $\mathcal{R}$ is obtained by substituting thus determined $\alpha_s^{(i)}$ and $(r_1^{(i)}, \dots, r_{i-1}^{(i)})$ for, in respective order, α_s and $(r_1, \dots, r_{i-1})$ in $\mathcal{R}^{(i)}$, Eq. (2.6).

It should be noted here that the above definition of $\mathcal{R}^{(i)}$ is not unique. Instead of (i) of the definition, one can adopt

$$\mathcal{R}^{(i)} = \alpha_s^l \left(\sum_{j=0}^{i-1} r_j \alpha_s^j + \sum_{j=i}^{\infty} r_j \alpha_s^j \right). \quad (2.9)$$

Here the coefficients $(r_i, r_{i+1}, \dots)$ are arbitrarily chosen fixed ones within the “reasonable” range. The original PMS is the special case, $r_i = r_{i+1} = \dots = 0$, of this

more general setup. The PMS criterion can also be applied to more general case, e.g.,

$$\mathcal{R} = \sum_{j=0}^{\infty} \alpha_s^{l+j} \sum_{k=0}^{n_j} (r_{jk} + \sqrt{\alpha_s} r'_{jk}) \ln^k \alpha_s, \quad (2.10)$$

where n_j is 0 or some integer.

2.2. Quark-gluon plasma pressure⁹⁾

As has been mentioned in §1, the form for the pressure p given in Ref. 9) consists of three pieces, $p = p_E + p_M + p_G$. We will apply the PMS criterion to p_E and p_{M+G} ($\equiv p_M + p_G$) separately. To define the border between the soft and hard energy regions, the scale parameter Λ_E is introduced, $O(g_s T) < O(\Lambda_E) < O(T)$.

The long-distance pressure p_{M+G} is given⁹⁾ as an expansion in powers of the effective EQCD parameters g_E and m_E :

$$p_{M+G} = \frac{2}{3\pi} T m_E^3 \tilde{p}_{M+G}, \quad (2.11)$$

$$\begin{aligned} \tilde{p}_{M+G} = & 1 - \frac{9}{4\pi} \frac{g_E^2}{m_E} \left(\frac{3}{4} + \ln \frac{\Lambda_E}{2m_E} \right) - \frac{110.2}{(4\pi)^2} \left(\frac{g_E^2}{m_E} \right)^2 \\ & + \frac{81}{(4\pi)^3} \left(\frac{g_E^2}{m_E} \right)^3 \left[4.44 \ln \frac{\Lambda_E}{2m_E} + 1.566 \ln \frac{\Lambda_E}{6g_E^2} - 1.392 + \delta_G \right] \\ & - \frac{5}{4\pi} (9 - n_f) \frac{\alpha_s^2}{m_E} T, \end{aligned} \quad (2.12)$$

where

$$m_E^2 = 4\pi(1 + n_f/6)\alpha_s T^2 \left[1 + \alpha_s \left(\beta_0 \ln \frac{\mu}{2\pi T} + \frac{1}{\pi} P_m(n_f) \right) \right], \quad (2.13)$$

$$g_E^2 = 4\pi\alpha_s T \left[1 + \alpha_s \left(\beta_0 \ln \frac{\mu}{2\pi T} + \frac{1}{\pi} P_g(n_f) \right) \right], \quad (2.14)$$

with

$$\begin{aligned} P_m(n_f) &= \frac{0.6124 - 0.4881n_f - 0.04280n_f^2}{1 + n_f/6}, \\ P_g(n_f) &= -0.3876 - 0.4235n_f. \end{aligned} \quad (2.15)$$

The dimensionless number δ_G in Eq. (2.12) is unknown. We will allow, as in Ref. 9), the following variation:

$$-5 < \delta_G < +5. \quad (2.16)$$

We refer⁶⁾ the expression (2.11) - (2.12) to as the untruncated form for p_{M+G} .

Expanding p_{M+G} , Eqs. (2.11) and (2.12), in powers of $\alpha_s^{1/2}$ and keeping up to and including $O(\alpha_s^3)$ contribution, we obtain the truncated form for p_{M+G} ,

$$p_{M+G}^{(2)} = p_{1,M+G}^{(2)} + p_{2,M+G}^{(2)}, \quad (2.17)$$

$$p_{1,M+G}^{(2)} = \alpha_s^{3/2} (r_{10} + r_{11}\alpha_s), \quad (2.18)$$

$$p_{2,M+G}^{(2)} = \alpha_s^2 (r_{20} + r_{21}\alpha_s). \quad (2.19)$$

We see from these expressions that, for p_{M+G} , μ is the only renormalization-scheme parameter and β_2 is not. The reason is as follows. On the one hand the highest-order term in $p_{M+G}^{(2)}$ is of $O(\alpha_s^3)$ (see Eq. (2.19)). On the other hand, a change in β_2 results in a change in $p_{M+G}^{(2)}$ of $O(\alpha_s^{7/2})$ ($< O(\alpha_s^3)$) (cf. Eq. (2.18) and the comment above after Eq. (2.5)). Thus, β_2 is not the renormalization-scheme parameter. When the $O(\alpha_s^{7/2})$ term of p_{M+G} is computed, β_2 becomes a renormalization-scheme parameter.

The short-distance pressure p_E is given⁹⁾ as an expansion in powers of the QCD couplant α_s :

$$p_E = p_{\text{SB}} \left(1 - \frac{15(1 + 5n_f/12)}{4(1 + 21n_f/32)} \tilde{p}_E \right), \quad (2.20)$$

$$p_{\text{SB}} = \frac{8\pi^2}{45} \left(1 + \frac{21}{32} n_f \right) T^4, \quad (2.21)$$

$$\begin{aligned} \tilde{p}_E = \frac{\alpha_s}{\pi} & \left[1 + \frac{\alpha_s}{\pi} \left(\pi\beta_0 \ln \frac{\mu}{2\pi T} - 36 \frac{1 + n_f/6}{1 + 5n_f/12} \left(\ln \frac{\Lambda_E}{2\pi T} + P_h(n_f) \right) \right) \right. \\ & + \left(\frac{\alpha_s}{\pi} \right)^2 \left((\pi\beta_0)^2 \ln^2 \frac{\mu}{2\pi T} + F(\Lambda_E/(2\pi T)) \right. \\ & \left. \left. + \left\{ \pi^2\beta_1 - 72\pi\beta_0 \frac{1 + n_f/6}{1 + 5n_f/12} \left(\ln \frac{\Lambda_E}{2\pi T} + P_h(n_f) \right) \right\} \ln \frac{\mu}{2\pi T} + \delta_E \right) \right], \end{aligned} \quad (2.22)$$

where

$$\begin{aligned} P_h(n_f) &= \frac{244.9 + 17.24n_f - 0.4150n_f^2}{135(1 + n_f/6)}, \\ F(\Lambda_E/2\pi T) &= \frac{478.4 + 35.14n_f + 4.081n_f^2}{1 + 5n_f/12} \left(\ln \frac{\Lambda_E}{2\pi T} + P_h(n_f) \right). \end{aligned} \quad (2.23)$$

The renormalization-scheme parameters for p_E is μ and β_2 . Eq. (2.22) includes the unknown number δ_E , which depends on the parameter β_2 . To see this dependence, we use the fact that the physical quantity p_E , Eq. (2.20), is independent of the parameters, μ and β_2 , up to and including $O(\alpha_s^3)$. Then from Eqs. (2.5) and (2.22), we find the relation between δ_E in the renormalization scheme with β_2 and its $\overline{\text{MS}}$ scheme counterpart $(\delta_E)_{\overline{\text{MS}}}$:

$$\delta_E = (\delta_E)_{\overline{\text{MS}}} - \frac{\pi^2}{\beta_0} [\beta_2 - (\beta_2)_{\overline{\text{MS}}}] . \quad (2.24)$$

For the unknown $(\delta_E)_{\overline{\text{MS}}}$, following Ref. 9), we will allow the following variation:

$$-|F(\Lambda_E/(2\pi T))| < (\delta_E)_{\overline{\text{MS}}} < +|F(\Lambda_E/(2\pi T))|. \quad (2.25)$$

While the pure number $(\delta_E)_{\overline{\text{MS}}}$ is independent of any scale, F will have only slight dependence on temperature T .

Finally, we note that, when the expression $p_{M+G}^{(2)}$, Eq. (2.17), is used for the long-distance pressure, the total pressure $p_{M+G}^{(2)} + p_E$ is independent⁹⁾ of the “separation

scale” Λ_E as it should be. It is worth remarking in passing that, if μ in $p_{M+G}^{(2)}$ is not equal to μ in p_E , this is not the case.

§3. Optimum approximants

It is found in Ref. 6) that various approximants, which include the optimal approximant, constructed using the untruncated form (2·11) - (2·12) for p_{M+G} are more reliable than those constructed using the truncated form (2·17) - (2·19). It is found in Ref. 9) that this is also the case for the approximants constructed on the basis of the Padé and Borel-Padé resummations of p_{M+G} . Keeping these observations and the remarks made above at the end of §2.1, we will also take more “reliable form” (2·11) - (2·12) for p_{M+G} , and apply the PMS criterion to it.

Before entering into our analysis, it is worth summarizing here the procedure and the results of applying the PMS criterion in Ref. 6).

1. The PMS criterion is applied to the total pressure $p = p_E + p_{M+G}$.
2. The 2-loop running couplant $\alpha_s^{(2)}(\mu)$ is used, so that the renormalization-scheme parameter is μ only.
3. Λ_E is chosen to be $\Lambda_E = \mu$.

The resultant optimal approximants are close to the lattice results for $T \gtrsim 0.7$ GeV. The three-loop optimal pressure (with three quark flavors) is $\sim 30\%$ larger than the lattice result at $T \simeq 0.3$ GeV, and the four-loop optimal pressure in pure-gluon QCD with appropriate choice for δ_E and δ_{M+G} is $\sim 20\%$ larger at $T \simeq 0.5$ GeV.

We will improve 1., 2., and 3. above as follows:

- 1'. The PMS criterion is applied to the short-distance pressure p_E and the long-distance pressure p_{M+G} separately.
- 2'. For p_E , the three-loop running couplant $\alpha_s^{(3)}(\mu)$ is used^{*)}, while, for p_{M+G} , the two-loop $\alpha_s^{(2)}(\mu)$ is used (see above after Eq. (2·19)).
- 3'. Λ_E is determined by requiring the minimal sensitivity of p on Λ_E (cf. Eq. (3·4) below).

In the following, the “long-distance” (“short-distance”) μ and α_s are denoted by μ_{M+G} and α_s^{M+G} (μ_E and α_s^E), respectively. The OPT equations for p_{M+G} read (see above after Eq. (2·19))

$$\mu_{M+G} \frac{\partial p_{M+G}}{\partial \mu_{M+G}} = 0, \quad \mu_{M+G} \frac{\partial \alpha_s^{M+G}}{\partial \mu_{M+G}} = \beta^{(2)}(\alpha_s^{M+G}), \quad (3·1)$$

where p_{M+G} is as in Eq. (2·11) with Eq. (2·12). The OPT equations for p_E read

$$\mu_E \frac{\partial p_E}{\partial \mu_E} = 0, \quad \frac{\partial p_E}{\partial \beta_2} = 0, \quad (3·2)$$

$$\mu_E \frac{\partial \alpha_s^E}{\partial \mu_E} = \beta^{(3)}(\alpha_s^E), \quad \frac{\partial \alpha_s^E}{\partial \beta_2} = \gamma^{(3)}(\alpha_s^E), \quad (3·3)$$

^{*)} As observed above in conjunction with Eq. (2·24), when the PMS criterion is applied to p_E (Eqs. (2·20) and (2·22)) in a consistent manner, the 3-loop running couplant should be used.

where p_E is as in Eq. (2·20) with Eqs. (2·21) - (2·23).

The scale parameter Λ_E is arbitrarily chosen in the range $O(g_s T) < O(\Lambda_E) < O(T)$, so that $p = p_{M+G} + p_E$ should not depend on Λ_E . As has been mentioned at the end of §2, this is the case provided that (α) the truncated expansion $p_{M+G}^{(2)}$ (in powers of $\alpha_s^{1/2}$) is used for p_{M+G} and (β) $\mu_E = \mu_{M+G}$. In our case, however, both (α) and (β) are not fulfilled, and then p *does* depend on Λ_E . To resolve this ambiguity, we introduce a criterion for determining Λ_E : Respecting the spirit of the PMS criterion, we employ the following determining equation for Λ_E ,

$$\frac{\partial p(T, \Lambda_E)}{\partial \Lambda_E} = 0. \quad (3.4)$$

Solving a set of equations (3.1) - (3.4), we obtain the optimized μ_{M+G} ($= \mu_L(T)$), α_s^{M+G} ($= \alpha_s^L(\mu_L(T))$), $\mu_E = \mu_S(T)$, $\beta_2 = \beta_{2S}(T)$, $\alpha_s^E = \alpha_s^S(\mu_S(T), \beta_{2S}(T))$ and $\Lambda_E(T)$ ($= \Lambda_E^{L/S}(T)$). Here “L” and “S” stand for “long-distance” and “short-distance”, respectively. Using thus determined quantities in Eqs. (2.11) and (2.20), we obtain the optimized pressure $p(T) = p_{M+G}(T) + p_E(T)$. Since $p = p_{M+G} + p_E$ is linear in Λ_E , when the μ_{M+G} , μ_E and β_2 are fixed to the respective optimized values, $p(\mu_L(T), \mu_S(T), \beta_{2S}(T), \Lambda_E; T)$ is independent of Λ_E .

§4. Results

We are concerned with the temperature range $0.3 \text{ GeV} \leq T \leq 3 \text{ GeV}$. Unless otherwise stated, we will take for the number of active (massless) quark flavors $n_f = 3$, which is of physical interest. For the QCD couplant $\alpha_s(\mu)$, we take the reference value $(\alpha_s(1.2 \text{ GeV}))_{\overline{\text{MS}}} = 0.403$. The form for the short-distance $\alpha_s^E(\mu_E, \beta_2)$ is obtained from Eq. (2.7), which is give in Appendix A. Following the procedure stated in §3, we obtain two solutions within the reasonable ranges for μ_{M+G} , μ_E and Λ_E . We refer the solution with larger (smaller) optimal approximant for p the solution I (II).

For the time being, we choose $\delta_G = (\delta_E)_{\overline{\text{MS}}} = 0$. Similar behaviors are observed for other values of δ_G and $(\delta_E)_{\overline{\text{MS}}}$.

In Fig. 1, we present the solutions I and II against T . The results are normalized by the Stefan-Boltzmann pressure p_{SB} (cf. Eq. (2.21)). Also shown are the optimized long-distance pressure p_{M+G} and the short-distance pressure p_E . The circles are the predictions of the lattice calculations¹¹⁾ (see Ref. 9) for details.) From the figures we see that i) for the solution II, $p_{M+G} < 0$ and ii) the solution I gives better agreement with the lattice result. From these observations, in the following, we adopt the solution I as the better candidate for the optimal approximant.

In Fig. 2 the optimized μ_{M+G} , $\mu_L(T)$, and μ_E , $\mu_S(T)$, are shown against T . The dot-dashed lines shows the traditional choice $\mu = 2\pi T$. The ratios $\mu_S(T)/T$ and $\mu_L(T)/T$ decrease monotonically with T : The ratio $\mu_S(T)/2\pi T$ is larger than 1 at $T = 0.3 \text{ GeV}$, $\simeq 1$ at $T = 0.5 \text{ GeV}$ and becomes smaller than 1 for $T > 0.5 \text{ GeV}$. $\mu_L(T)$ is smaller than $\mu_S(T)$ and $\mu_L(T)/\mu_S(T) \simeq 0.31 \sim 0.36$ in the range shown in the figure. This is in accord with an intuition: The scale μ_E that governs the physics

in the hard region, being of $O(T)$, is larger than the scale μ_{G+M} in the soft region.

Fig. 3 depicts the optimized β_2 , $\beta_{2S}(T)$, against T . The dashed line shows $\beta_2 = (\beta_2)_{\overline{\text{MS}}}$. $\beta_{2S}(T)$ monotonically decreases with T crossing the $\overline{\text{MS}}$ value at $T \simeq 0.5$ GeV. We note that Eq. (2.5) tells us that increase (decrease) in β_2 acts as increasing (decreasing) α_s , the same effect as arising from decreasing (increasing) μ . Then, we suspect that, if we fix β_2 to $(\beta_2)_{\overline{\text{MS}}}$ and vary only μ , we obtain much smaller (larger) optimized μ_E for $T \lesssim 0.5$ ($T \gtrsim 0.5$) GeV. Thus, we expect that the optimized μ_E tends to $2\pi T$. It is amazing to note that, at $T \sim 0.5$ GeV, $\mu_S(T)$ and $\beta_{2S}(T)$ are close to the traditional choice $\mu = 2\pi T$ and the $\overline{\text{MS}}$ value, respectively.

In Fig. 4, we show the optimized Λ_E , $\Lambda_E^{L/S}(T)$, and $m_E(T)$ against T . For reference $\mu_L(T)$ in Fig. 2 is also shown again. We see that $\Lambda_E^{L/S}(T) \sim m_E(T) \sim \mu_L(T) (< \mu_S(T))$. This is not in accord with an intuitive expectation $\mu_{M+G} < \Lambda_E < \mu_E$. However, as seen at the end of §3, when μ_{M+G} , μ_E , and β_2 are fixed to their respective optimized values, the pressure p does not depend on Λ_E .

In Fig. 5, we show the pressure p at $T = 0.5$ GeV near the optimized values of $\alpha_s^E \simeq \alpha_s^S(T)$, $\alpha_s^{M+G} \simeq \alpha_s^L(T)$, $\beta_2 \simeq \beta_{2S}(T)$ and $\Lambda_E \simeq \Lambda_E^{L/S}(T)$. In Fig. 5 (a), we show the behavior of p against α_s^S , when other quantities, α_s^{M+G} , β_2 and Λ_E are fixed to the respective optimized values. The arrow indicates the optimized value of α_s^E . Similarly, Figs. 5 (b) and (c) show the dependencies of p on α_s^{M+G} and on β_2 , respectively. From Fig. 5(c), we see that p depends on β_2 only weakly. As has been repeatedly pointed out, p does not depend on Λ_E , when α_s^E , α_s^{M+G} and β_2 are fixed to the respective optimized values.

In Fig. 6, we present the optimal approximants for p , when the unknown constants $(\delta_E)_{\overline{\text{MS}}}$ and δ_G are varied. Within the ranges, (2.16) and (2.25), of the unknown constants, $(\delta_G, (\delta_E)_{\overline{\text{MS}}}) = (+5, -F)$ yields the maximum p/p_{SB} , while $(\delta_G, (\delta_E)_{\overline{\text{MS}}}) = (-5, +F)$ yields the minimum p/p_{SB} . For comparison, the $\overline{\text{MS}}$ results are also shown^{*)}. From the figure, we see as a bonus that the resultant optimal approximants depend rather weakly on δ_G and $(\delta_E)_{\overline{\text{MS}}}$. Our results are substantially improved when compared to the $\overline{\text{MS}}$ results. Taking into account that the lattice results should be taken within an error of 10 – 15 %, our results are well compatible with the lattice results.

Finally, in Fig. 7, we depicts the optimal approximants (solutions I) for the case of pure glue QCD ($n_f = 0$). For comparison, the solution II for $\delta_G = (\delta_E)_{\overline{\text{MS}}} = 0$ is also shown. Same comments as for the $n_f = 3$ case applies here. For $\delta_G = (\delta_E)_{\overline{\text{MS}}} = 0$, the $\overline{\text{MS}}$ result accidentally coincides with the solution II with high precision. Although the results are substantially improved when compared to the $\overline{\text{MS}}$ results (except for the case of $(\delta_G, (\delta_E)_{\overline{\text{MS}}}) = (+5, -F)$), agreement with the lattice results

^{*)} p_{M+G} (Eq. (2.12)) contains the logarithmic factors $\ln m_E$ and $\ln g_E^2$, for which we have expanded as $\ln \alpha_s + c_1 + c_2 \alpha_s + \dots$ with respective constants $c_1, c_2, \dots$. Then, for p_{M+G} , we have used the truncated forms (2.17) - (2.19) and then p ($= p_{M+G} + p_E$) is independent of Λ_E . For the $\beta(\alpha_s)$ function, we have used the three-loop form. For the scale parameter μ , we have used the conventional one $\mu = 2\pi T$. The unknown constant $(\delta_E)_{\overline{\text{MS}}}$ in Eq. (2.22), we vary in the range (2.25). For Λ_E in $F(\Lambda_E/2\pi T)$ there, we have used Eq. (2.23) with $\Lambda_E = \sqrt{2\pi T m_E(T)}$. For μ_{M+G} and α_s^{M+G} in $m_E(\mu_{M+G}, \alpha_s^{M+G}, T)$, we have substituted the respective optimized values.

are less than that for the realistic case of $n_f = 3$.

§5. Discussion and Conclusion

The QCD pressure p has been computed within the improved perturbation theory using the $\overline{\text{MS}}$ renormalization scheme, up to and including the $O(g_s^6 \ln(1/g_s))$ and part of the $O(g_s^6)$ terms. The truncated perturbation series for p does not reconcile with the relatively low T lattice results. As has been mentioned in §1, various approaches are proposed for improving the perturbative pressure p . Each approach performs a sort of resummation on the basis of respective philosophy. In this paper we add one more to them.

We have applied the Principles of Minimal Sensitivity (PMS) criterion to p . This is not new, but in Ref. 6), among other things, the PMS criterion has already been applied to p . However, as has been mentioned in §3, application is not made in a consistent manner. Meanwhile, an important observation is made in Ref. 9): The contributions to p from the hard and soft energy regions, p_E and p_{M+G} , are separately physical quantities. On the basis of this fact, we have applied the PMS criterion to p_E and p_{M+G} separately. For determining the factorization scale Λ_E , which separates hard and soft scales, we also adopt the minimal sensitivity criterion.

We have found that the optimal approximants are significantly improved when compared to the $\overline{\text{MS}}$ results. Especially, for the realistic case of $n_f = 3$, agreement between our approximants and the lattice results is satisfactory. Moreover, we have found that, when compared to the $\overline{\text{MS}}$ results, the dependence of the approximants on the unknown terms of $O(g_s^6)$ is rather weak. On the other hand, for the case of $n_f = 0$, although the results are substantially improved, agreement with the lattice results are poorer than that for the $n_f = 3$ case.

Further improvement may be expected if we perform the Padé type resummation starting from the optimal approximants obtained here, which is outside the scope of this paper.

Acknowledgments

We thank A. Nakamura for instructions of lattice calculations. The authors thank the useful discussion at the Workshop on Thermal Field Theories and their Applications, held at the Yukawa Institute for Theoretical Physics, Kyoto, Japan, 9 - 11 August, 2004. A. N. is supported in part by the Grant-in-Aid for Scientific Research [(C)(2) No. 17540271] from the Ministry of Education, Culture, Sports, Science and Technology, Japan, No.(C)(2)-17540271.

Appendix A: Running Couplant

The relation between the three-loop running couplants $\alpha_s(\mu_0, (\beta_2)_{\overline{\text{MS}}})$ and $\alpha_s(\mu, \beta_2)$ is obtained by integrating the following two equations (cf. Eq. (2.7)),

$$\mu \frac{\partial \alpha_s}{\partial \mu} = \beta^{(3)}(\alpha_s), \quad (\text{A.1})$$

$$\frac{\partial \alpha_s}{\partial \beta_2} = \gamma_2^{(3)}(\alpha_s), \quad (\text{A.2})$$

along a path $\mathcal{C}$ in a $(\ln \mu, \beta_2)$ -plane, which starts from the point $(\ln \mu_0, (\beta_2)_{\overline{\text{MS}}})$ and arrives at the point $(\ln \mu, \beta_2)$. Because of the integrability condition $\partial \beta^{(3)}(\alpha_s) / \partial \beta_2 = \partial \gamma_2^{(3)}(\alpha_s) / \partial \ln \mu$, one can use any path $\mathcal{C}$. For convenience, we choose $\mathcal{C} = \mathcal{C}_1 \oplus \mathcal{C}_2 \oplus \mathcal{C}_3$ with $\mathcal{C}_1 = (\ln \mu_0, (\beta_2)_{\overline{\text{MS}}}) \rightarrow (\ln \mu_M, (\beta_2)_{\overline{\text{MS}}})$, $\mathcal{C}_2 = (\ln \mu_M, (\beta_2)_{\overline{\text{MS}}}) \rightarrow (\ln \mu_M, \beta_2)$ and $\mathcal{C}_3 = (\ln \mu_M, \beta_2) \rightarrow (\ln \mu, \beta_2)$. At the final stage, we take the limit $\mu_M \rightarrow +\infty$.

Integrating Eq. (A.1) along the path $\mathcal{C}_1$, we obtain

$$\begin{aligned} \beta_0 \ln \left(\frac{\mu_M}{\mu_0} \right) &= \frac{1}{\alpha_{sM}} - \frac{1}{\alpha_{s0}} + \frac{\beta_1}{\beta_0} \ln \left(\frac{\alpha_{sM}}{\alpha_{s0}} \right) - \frac{\beta_1}{2\beta_0} \ln \left(\frac{\beta_0 + \beta_1 \alpha_{sM} + (\beta_2)_{\overline{\text{MS}}} \alpha_{sM}^2}{\beta_0 + \beta_1 \alpha_{s0} + (\beta_2)_{\overline{\text{MS}}} \alpha_{s0}^2} \right) \\ &\quad - \frac{\beta_1^2 - 2\beta_0 (\beta_2)_{\overline{\text{MS}}}}{2\beta_0^2} [G(\alpha_{sM}, (\beta_2)_{\overline{\text{MS}}}) - G(\alpha_{s0}, (\beta_2)_{\overline{\text{MS}}})], \end{aligned} \quad (\text{A.3})$$

where $\alpha_{s0} = \alpha_s(\mu_0, (\beta_2)_{\overline{\text{MS}}})$ and $\alpha_{sM} = \alpha_s(\mu_M, (\beta_2)_{\overline{\text{MS}}})$ and

$$G(x, y) = \frac{2\beta_0}{\sqrt{4\beta_0 y - \beta_1^2}} \arctan \left(\frac{2xy + \beta_1}{\sqrt{4\beta_0 y - \beta_1^2}} \right).$$

Integration of Eq. (A.1) along the path $\mathcal{C}_3$ yields a similar equation for $\beta_0 \ln(\mu/\mu_M)$. Adding this to Eq. (A.3), we obtain the formula for $\beta_0 \ln(\mu/\mu_0)$, which includes, among others, $\alpha_s(\mu_M, (\beta_2)_{\overline{\text{MS}}})$ and $\alpha_s(\mu_M, \beta_2)$. From Eq. (A.2), we see that $\alpha_s(\mu_M, (\beta_2)_{\overline{\text{MS}}}) = \alpha_s(\mu_M, \beta_2) + O(\alpha_s^3)$. Substituting this for $\alpha_s(\mu_M, (\beta_2)_{\overline{\text{MS}}})$ in the formula obtained above and taking the limit $\mu_M \rightarrow +\infty$ ($\alpha_s(\mu_M, \beta_2) \rightarrow 0$), we see that the $O(\alpha_s^3)$ term does not survive.

Thus, we finally obtain

$$\begin{aligned} \beta_0 \ln \left(\frac{\mu}{\mu_0} \right) &= \frac{1}{\alpha_s} - \frac{1}{\alpha_{s0}} + \frac{\beta_1}{\beta_0} \ln \left(\frac{\alpha_s}{\alpha_{s0}} \right) - \frac{\beta_1}{2\beta_0} \ln \left(\frac{\beta_0 + \beta_1 \alpha_s + \beta_2 \alpha_s^2}{\beta_0 + \beta_1 \alpha_{s0} + (\beta_2)_{\overline{\text{MS}}} \alpha_{s0}^2} \right) \\ &\quad - \frac{\beta_1^2 - 2\beta_0 (\beta_2)_{\overline{\text{MS}}}}{2\beta_0^2} [G(0, (\beta_2)_{\overline{\text{MS}}}) - G(\alpha_{s0}, (\beta_2)_{\overline{\text{MS}}})] \\ &\quad - \frac{\beta_1^2 - 2\beta_0 \beta_2}{2\beta_0^2} [G(\alpha_s, \beta_2) - G(0, \beta_2)], \end{aligned} \quad (\text{A.4})$$

where $\alpha_s = \alpha_s(\mu, \beta_2)$.

References

- 1) P. Arnold and C. Zhai, Phys. Rev. D **50**, 7603 (1994); *ibid.* **51**, 1906 (1995); C. Zhai and B. Kastening, *ibid.* **52**, 7232 (1995); E. Braaten and A. Nieto, Phys. Rev. Lett. **76**, 1417 (1996); Phys. Rev. D **53**, 3421 (1996).

- 2) K. Kajantie, M. Laine, K. Rummukainen, and Y. Schröder, Phys. Rev. D **67**, 105008 (2003).
- 3) A. Peshier, B. Kämpfer, O. P. Pavlenko, and G. Soff, Phys. Rev. D **54**, 2399 (1996); P. Lévai and U. Heinz, Phys. Rev. C **57**, 1879 (1998).
- 4) J.-P. Blaizot, E. Iancu, and A. Rebhan, Phys. Lett. **B470**, 181 (1999); Phys. Rev. Lett. **83**, 2906 (1999); A. Peshier, Phys. Rev. D **63**, 105004 (2001).
- 5) F. Karsch, A. Patkós, and P. Petreczky, Phys. Lett. **B401**, 69 (1997); J. O. Andersen and M. Strickland, Phys. Rev. D **64**, 105012 (2001).
- 6) J.-P. Blaizot, E. Iancu, and A. Rebhan, Phys. Rev. D **68**, 025011 (2003).
- 7) J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. D **61**, 014017 (2000); *ibid.* **61**, 074016 (2000); J. O. Andersen, E. Petitgirard, and M. Strickland, *ibid.* **70**, 045001 (2004).
- 8) B. Kastening, Phys. Rev. D **56**, 8107 (1997); T. Hatsuda, *ibid.* **56**, 8111 (1997).
- 9) G. Cvetič and R. Kögerler, Phys. Rev. D **70**, 114016 (2004); *ibid.* **66**, 105009 (2002).
- 10) P. M. Stevenson, Phys. Rev. D **23**, 2916 (1981).
- 11) F. Karsch, E. Laermann, and A. Peikert, Phys. Lett. **B478**, 447 (2000); F. Karsch, Nucl. Phys. **A698**, 199 (2002).




Fig. 1. The solutions I and II for the optimal approximant, p/p_{SB} , as functions of T , when $\delta_G = (\delta_E)_{\overline{\text{MS}}} = 0$. (p_{SB} is the Stefan-Boltzmann pressure.) The optimized long-distance pressure p_{M+G} and the short-distance pressure p_E are also separately shown. The circles are the predictions of the lattice calculations.




Fig. 2. The optimized μ_{M+G} , $\mu_L(T)$, and μ_E , $\mu_S(T)$, for $\delta_G = (\delta_E)_{\overline{\text{MS}}} = 0$ as functions of T . The dot-dashed lines shows the traditional choice $\mu = 2\pi T$.




Fig. 3. The optimized β_2 , $\beta_{2S}(T)$, as a function of T . The dashed line shows $\beta_2 = (\beta_2)_{\overline{\text{MS}}}$.




Fig. 4. The optimized Λ_E , $\Lambda_E^{L/S}(T)$, and m_E as functions of T . For reference, the optimized μ_{M+G} , $\mu_L(T)$, is also shown.




Fig. 5. The pressure p near the optimized values of α_s^E , α_s^{M+G} and β_2 , when $T = 0.5$ GeV. The arrows in Figs. (a), (b) and (c) indicate the optimized values of α_s^E , α_s^{M+G} and β_2 , respectively. Details are given in the text.




Fig. 6. Optimal approximants (the solutions I) for p/p_{SB} as functions of T for various values of δ_G and $(\delta_E)_{\overline{\text{MS}}}$. Within the ranges, (2.16) and (2.25), of $(\delta_G, (\delta_E)_{\overline{\text{MS}}})$, $(\delta_G, (\delta_E)_{\overline{\text{MS}}}) = (+5, -F)$, yields the maximum p/p_{SB} (dot-dashed line) while $(\delta_G, (\delta_E)_{\overline{\text{MS}}}) = (-5, +F)$ yields the minimum one (solid line). For comparison, the lattice results and the $\overline{\text{MS}}$ results are also shown. Displayed $\overline{\text{MS}}$ results are the maximum one [$(\delta_G, (\delta_E)_{\overline{\text{MS}}}) = (+5, -F)$], within the ranges (2.16) and (2.25), the minimum one [$(\delta_G, (\delta_E)_{\overline{\text{MS}}}) = (-5, +F)$] and the one with $\delta_G = (\delta_E)_{\overline{\text{MS}}} = 0$.




Fig. 7. Analogous to Fig. 6 but with $n_f = 0$. The solution II for $\delta_G = (\delta_E)_{\overline{\text{MS}}} = 0$ is also shown.